

## Prokaryotic toxin-antitoxin systems — the role in bacterial physiology and application in molecular biology

Michał Bukowski, Anna Rojowska and Benedykt Władyka✉

Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Kraków, Poland

**Bacteria have developed multiple complex mechanisms ensuring an adequate response to environmental changes. In this context, bacterial cell division and growth are subject to strict control to ensure metabolic balance and cell survival. A plethora of studies cast light on toxin-antitoxin (TA) systems as metabolism regulators acting in response to environmental stress conditions. Many of those studies suggest direct relations between the TA systems and the pathogenic potential or antibiotic resistance of relevant bacteria. Other studies point out that TA systems play a significant role in ensuring stability of mobile genetic material. The evolutionary origin and relations between various TA systems are still a subject of a debate. The impact of toxin-antitoxin systems on bacteria physiology prompted their application in molecular biology as tools allowing cloning of some hard-to-maintain genes, plasmid maintenance and production of recombinant proteins.**

**Keywords:** antibiotic resistance, bacteria physiology, environmental stress conditions, toxin-antitoxin systems

**Received:** 16 March, 2010; **revised:** 24 January, 2011; **accepted:** 08 March, 2011; **available on-line:** 11 March, 2011

### INTRODUCTION

Toxin-antitoxin systems emerged in research in mid 80's. A detailed insight into their functions and mechanisms of action has been gained in the last two decades and brought several interesting conclusions as to the importance of such systems for bacterial physiology. The term "toxin-antitoxin system", usually abbreviated as "TA system", comprises a functional element consisting, in most cases, of a biologically active protein molecule and a corresponding inhibitor, whose nature and inhibitory mechanism depend on the system's class affiliation. Components of such systems are encoded within polycistronic operons, often with partially overlapping open reading frames. The systems are widespread among *Bacteria* as well as *Archaea* (Mittenhuber, 1999; Gerdes, 2000; Pandey & Gerdes, 2005; Makarova *et al.*, 2009) and evolved to carry out diverse functions. However, their common feature is an enzymatic activity detrimental for the cell metabolism. Such toxic activity has been demonstrated to switch bacterial cells over to a dormant state, leading to cell death during prolonged exposure. In most cases various stress stimuli are responsible for TA system activation. The signalling pathway in such instances is often related to other stress-induced response pathways. Moreover, it is well documented that in some cases the activity of TA systems stabilizes mobile genetic elements, therefore comprising an important mechanism

of plasmids maintenance. In the light of the increasing multi-drug resistance among virulent strains, reports on the potential relation between TA systems and modulation of pathogen–host interactions seem to be of utmost importance.

### CLASSIFICATION OF TOXIN-ANTITOXIN SYSTEMS

The biological activity of a toxin comprising a component of a TA systems is usually (but not always) that of an endoribonuclease. Bioinformatic analysis of multiple available sequences of bacterial genetic elements points to multiple novel, putative TA *loci* and suggests that many of known TA systems, bacterial as well as archaeal, are evolutionarily related (Anantharaman & Aravind, 2003; Hayes & Sauer, 2003; Gerdes *et al.*, 2005; Sevin & Barloy-Hubler, 2007; Makarova *et al.*, 2009; Weaver *et al.*, 2009; Arbing *et al.*, 2010). The classification of TA systems is based on the mechanism of inhibition of the toxin as well as on operon autoregulatory functions. Initially two classes of TA systems were identified (Gerdes & Wagner, 2007), but subsequent discoveries extended the classification to three classes (Blower *et al.*, 2009). Recent studies suggest the existence of yet another type, namely a three-component TA system (Hallez *et al.*, 2010). As immediately visible from the above discussion the field is in a constant and dynamic growth and one may expect that many interesting findings are likely to emerge in the following years.

Class I includes systems in which the antitoxin is an antisense RNA forming duplexes with the toxin mRNA. This leads to inhibition of translation in a process known as RNA interference. Examples of such systems are chromosomally located operons found in *Escherichia coli*, namely *tisAB* (Vogel *et al.*, 2004) and *symER* (Kawano *et al.*, 2007), as well as plasmid loci *parB* (Gerdes *et al.*, 1986) of *E. coli* and *par* of *Enterococcus faecalis* (Greenfield *et al.*, 2000; Weaver *et al.*, 2009) and a homologous plasmid operon of *Staphylococcus aureus* (Jensen *et al.*, 2010). Among the mentioned systems toxins have multiple different roles. For example the SymE toxin is an mRNA interferase encoded in the *symER* operon. The toxin binds ribosomes to exert its activity (Kawano *et al.*, 2007). The TisB toxin, which is encoded in the *tisAB* operon (Vogel *et al.*, 2004) decreases the proton-motric force across the bacterial cell membrane and cause subsequent drop in ATP production, which leads

✉ e-mail: wladykab@interia.pl

**Abbreviations:** TA system, toxin-antitoxin system; ppGpp, 3',5'-guanosine bisphosphate; NMR, nuclear magnetic resonance; SPP system, single protein production system

to metabolic dormancy (Unoson & Wagner, 2008). Hok toxin, encoded in the *parB* operon, irreversibly damages the cell membrane (Gerdes *et al.*, 1986). In the latter case the regulation of the toxin level is indirect. RNA interference suppresses expression of the gene *mok*, which is a regulator of *hok* gene transcription (Thisted & Gerdes, 1992).

Class II encompasses a wide range of TA systems. Antitoxins of this class are proteins. The biological activities exhibited by the toxins include transcription inhibition by targeting gyrase function and interference with translation through an mRNA interferase activity, which may or may not rely on ribosome binding. The endoribonucleolytic activity of mRNA interferases is often sequence specific. Table 1 gives a short overview of the class II TA systems and their characteristics.

Class III comprises a single member only. This system is encoded in the *toxIN* operon of *Erwinia carotovora*, a plant pathogen. In this case inhibition of ToxN toxin activity is driven by RNA molecules directly interacting with the toxin molecules (Blower *et al.*, 2009; Fineran *et al.*, 2009).

#### RELATIONS AND STRUCTURAL SIMILARITIES AMONG CLASS II TA SYSTEMS


The evolutionary relationship among class II TA systems is a subject of an open debate. Attention is mainly focused on toxins since there is a substantial sequence and structural variety among the antitoxins. Ten TA families of class II have been described so far (Pandey & Gerdes, 2005; Jorgensen *et al.*, 2009; Van Melderden & Saavedra De Bast, 2009) and for three of them, *relBE*, *parDE* and *bigBA*, a phylogenetic relationship based on sequence similarities has been proposed (Anantharaman & Aravind, 2003; Tsilibaris *et al.*, 2007). Strikingly, the toxin of the *parDE* system is a gyrase inhibitor in contrast to the toxins of the *relBE* and *bigBA* systems, which are mRNA interferases. A broader analysis of this

issue leads to other interesting conclusions. There is no evidence for an evolutionary relation between the *cdAB* and *parDE* systems (Anantharaman & Aravind, 2003) although the toxin of the *cdAB* system is also a gyrase inhibitor. However, there is a significant structural similarity between the toxins of the *cdAB* and *kis/kid* (*parD*) systems (Diago-Navarro *et al.*, 2010), which, similarly to the *parDE* and *relBE* or *bigBA* systems, are a gyrase inhibitor and an mRNA interferase, respectively. Other reports point to a structural similarity among the toxins of the *ygiUT* (*mqsRA*), *relBE* and *yefM-yoeB* systems as well as RNase Sa of *Streptomyces aureofaciens* (Brown *et al.*, 2009).

Not only among RelE homologues is a similarity with RNase Sa noticeable. Toxins of the *cdAB* and *kis/kid* or *mazEF* (*cbpAK*) systems are also structurally similar. This similarity is related to the presence of a  $\beta$ -sheet core in these molecules (Fig. 1). However, this  $\beta$ -sheet core structure is most likely related to the ability to form dimers (Miller, 1989) rather than reflects evolutionary or functional relationships. Structural analysis of mRNA interferases and comparative studies allow the deduction of the mechanism of their endoribonucleolytic activity (Agarwal *et al.*, 2009; Brown *et al.*, 2009; Diago-Navarro *et al.*, 2010). Tracing evolutionary relations among the TA systems is difficult because of the fast specialisation of TA system components (Arbing *et al.*, 2010). It has been reported that the toxin of the *phd/doc* system is similar to a virulence factor toxic to eukaryotic host cells (Arbing *et al.*, 2010). Another example is the sequence similarity of toxins of the *symER* and *phd/doc* systems to antitoxins of other TA systems — *yefM-yoeB* (Arbing *et al.*, 2010) and *mazEF* (Kawano *et al.*, 2007), respectively.

#### REGULATION OF CLASS II TA SYSTEM ACTIVITY

In operons of class II TA systems an antitoxin gene is usually, but not always, located upstream a gene for a toxin. The order is reversed for example in the *bigBA*,



**Figure 1. Structural similarities among toxins belonging to different families**

(A) *ccdBA* and *mazEF* (Diago-Navarro *et al.*, 2010); (B) *relBE* and RNase Sa of *Streptomyces aureofaciens* (Brown *et al.*, 2009). In fact,  $\beta$ -sheet core (red) structure is similar among all these toxins. Models prepared with PyMOL ver. 1.1r2pre (DeLano WL, 2002). Structures' PDB IDs — CcdB: 1VUB; Kid: 1M1F; MazF: 1UB4; RelE: 2KC8; YoeB: 2A6Q; MqsR: 3HI2; RNase Sa: 1RSN.

Table 1. Ten families of class II TA systems and data about well-researched members


Family	Operon	Toxin	Antitoxin	Source organism/location	Activity	Mechanism of toxicity
ccdAB	ccdAB	CcdB	CcdA	<i>Escherichia coli</i> /plasmid <sup>1</sup>	gyrase inhibitor <sup>2</sup>	transcription inhibition <sup>2</sup>
parDE	parDE	ParE	ParD	<i>Escherichia coli</i> /plasmid <sup>3</sup>	gyrase inhibitor <sup>4</sup>	transcription inhibition <sup>4</sup>
phd/doc	phd/doc	Phd	Doc	prophage P1 <sup>5</sup>	binding ribosome 30S subunit <sup>6</sup>	translation inhibition <sup>6</sup>
mazEF	mazEF (chpAK)	MazF (ChpK)	MazE (ChpA)	<i>Escherichia coli</i> /chromosome <sup>7</sup>	endoribonuclease <sup>8</sup>	translation inhibition <sup>8</sup>
	kis/kid (parD)	Kid	Kis	<i>Escherichia coli</i> /plasmid <sup>9</sup>	endoribonuclease <sup>10</sup>	translation inhibition <sup>10</sup>
	pemIK	PemK	PemI	<i>Escherichia coli</i> /plasmid <sup>11</sup>	endoribonuclease <sup>12</sup>	translation inhibition <sup>12</sup>
	chpBIK	ChpBK	ChpBI	<i>Escherichia coli</i> /chromosome <sup>13</sup>	endoribonuclease <sup>14</sup>	translation inhibition <sup>14</sup>
	mazEF-mt1 – mazE-mt7	MazF <sub>Sa</sub>	MazE-mt1 – MazE-mt7	<i>Mycobacterium tuberculosis</i> /chromosome <sup>15</sup>	MazF-mt-1,3,6,7 – endoribonuclease, others not researched <sup>15</sup>	MazF-mt-1,3,6,7 – translation inhibition, others not researched <sup>15</sup>
	mazEF <sub>Sa</sub>	MazF <sub>Sa</sub>	MazE <sub>Sa</sub>	<i>Staphylococcus aureus</i> /chromosome <sup>16</sup>	endoribonuclease <sup>16</sup>	translation inhibition <sup>16</sup>
	pemIK <sub>Sa</sub>	PemK <sub>Sa</sub>	PemI <sub>Sa</sub>	<i>Staphylococcus aureus</i> /plasmid <sup>17</sup>	endoribonuclease <sup>18</sup>	unknown
	relBE	RelE	RelB	<i>Escherichia coli</i> /chromosome <sup>19</sup>	endoribonuclease, ribosome-binding <sup>20</sup>	translation inhibition <sup>20</sup>
	yefM-yoeB	YoeB	YefM	<i>Escherichia coli</i> /chromosome <sup>21</sup>	endoribonuclease, ribosome-binding <sup>22</sup>	translation inhibition <sup>22</sup>
	yafNO	YafO	YafN	<i>Escherichia coli</i> /chromosome <sup>23</sup>	endoribonuclease, ribosome-binding <sup>23</sup>	translation inhibition <sup>23</sup>
ygjNM	YgjN	YgjM	<i>Escherichia coli</i> /chromosome <sup>23</sup>	endoribonuclease, ribosome-binding <sup>23</sup>	translation inhibition <sup>23</sup>	
ygiUT (mqsRA)	YgiU (MqsR)	YgiT (MqsA)	<i>Escherichia coli</i> /chromosome <sup>23</sup>	endoribonuclease <sup>23</sup>	translation inhibition <sup>23</sup>	
dinJ-yafQ	YafQ	DinJ	<i>Escherichia coli</i> /chromosome <sup>23</sup>	endoribonuclease, ribosome-binding <sup>23</sup>	translation inhibition <sup>23</sup>	
higBA	higB	HigA	<i>Vibrio cholerae</i> /chromosome <sup>24</sup>	endoribonuclease, ribosome-binding <sup>25</sup>	translation inhibition <sup>25</sup>	
vapBC	vapC	VapB	<i>Mycobacterium smegmatis</i> /chromosome <sup>26</sup>	endoribonuclease <sup>27</sup>	translation inhibition <sup>27</sup>	
ζε	ζ	ε	<i>Streptococcus pyogenes</i> /plasmid <sup>28</sup>	phosphotransferase <sup>29</sup>	unknown	
hipBA	hipA	HipB	<i>Escherichia coli</i> /chromosome <sup>30</sup>	Ser/Thr kinase (target: EF-Tu) <sup>31</sup>	translation inhibition <sup>32</sup>	
hicAB	hicA (ydcQ)	HicB (YdcQ)	<i>Escherichia coli</i> /chromosome <sup>33</sup>	endoribonuclease <sup>34</sup>	translation inhibition <sup>34</sup>	

<sup>1</sup>(Ogura & Hiraga, 1983); <sup>2</sup>(Miki et al., 1992); <sup>3</sup>(Saurugger, 1986); <sup>4</sup>(Jiang et al., 2002); <sup>5</sup>(Lehnherr et al., 1993; Magnuson & Yarmolinsky, 1998; Gazit & Sauer, 1999); <sup>6</sup>(Liu et al., 2008); <sup>7</sup>(Masuda et al., 1993); <sup>8</sup>(Munoz-Gomez et al., 2004); <sup>9</sup>(Bravo et al., 1987; Bravo et al., 1988); <sup>10</sup>(Zhang et al., 2003); <sup>11</sup>(Tsuchimoto et al., 1988); <sup>12</sup>(Zhang et al., 2004); <sup>13</sup>(Masuda et al., 1993); <sup>14</sup>(Zhang et al., 2005); <sup>15</sup>(Zhu et al., 2006); <sup>16</sup>(Fu et al., 2007; 2009; 2009; 2009; 2009); <sup>17</sup>(Lowder et al., 2009); <sup>18</sup>(Bukowski et al., 2010); <sup>19</sup>(Lavelle, 1965; Diderichsen et al., 1977; Bech et al., 1985; Mosteller, 1978); <sup>20</sup>(Galvani et al., 2001; Pedersen et al., 2003); <sup>21</sup>(Christensen et al., 2004); <sup>22</sup>(Christensen-Dalsgaard & Gerdes, 2008; Zhang & Inouye, 2009); <sup>23</sup>(Yamaguchi et al., 2010); <sup>24</sup>(Buddle et al., 2007); <sup>25</sup>(Christensen-Dalsgaard & Gerdes, 2006); <sup>26</sup>(Arcus et al., 2005); <sup>27</sup>(Daines et al., 2007; Robson et al., 2009); <sup>28</sup>(Camacho et al., 2002; Lloy et al., 2002); <sup>29</sup>(Meinhart et al., 2003); <sup>30</sup>(Black et al., 1991; 1994; 1994; Korch & Hill, 2006); <sup>31</sup>(Correia et al., 2006); <sup>32</sup>(Schumacher et al., 2009); <sup>33</sup>(Makarova et al., 2006); <sup>34</sup>(Jorgensen et al., 2009).

*bicAB* and *ygiUT* systems. Binding of toxin-antitoxin complexes to promoter sites is the most common way of direct transcription regulation of TA operons (Fig. 2). Single components also bind the promoters but with a low affinity (Kedzierska *et al.*, 2007; Li *et al.*, 2008) when compared to the toxin-antitoxin oligomers which bind to palindromic sequences within the promoters, which process is enhanced cooperatively (Tsuchimoto & Ohtsubo, 1993; Black *et al.*, 1994; Magnuson *et al.*, 1996; Magnuson & Yarmolinsky, 1998; Marianovsky *et al.*, 2001; Bailey & Hayes, 2009). Moreover, apart from the described primary palindromes, promoter of the *mazEF* operon contains alternate palindromes.

Binding to the latter by a toxin-antitoxin complex manifests in a decrease in the transcription efficiency of the operon (Marianovsky *et al.*, 2001). An exception to the above rule is the prophage P1 zeta-epsilon system ( $\zeta\epsilon$ ) where the antitoxin serves only as an inhibitor of toxin activity and an additional expression regulator  $\omega$  is present (de la Hoz *et al.*, 2000), which is similar to recently reported three-component systems homologous to *parDE*, namely *paaR1-paaA1-parE1* and *paaR2-paaA2-parE2* (Hallez *et al.*, 2010). Such a way of controlling the cellular levels of TA system components combined with high proteolysis susceptibility of the antitoxin provides the way of tight and environmentally switchable regulation. The instability of the antitoxin in a TA system is a crucial step in the system activation. It is suggested that disordered C-terminal regions of the antitoxin are target for ATP-dependent serine proteases (Kamada *et al.*, 2003). These members of chaperone family are responsible for degradation of misfolded proteins as well as components of signalling pathways (Gottesman, 1996). However, the antitoxin YgiT (MqsA) of the *ygiUT* (*mqsR4*) system is structured throughout its entire sequence, both free and toxin-bound state (Brown *et al.*, 2009). The activity of ATP-dependent proteases stays in a specific relation with the activity of TA systems. In all documented cases only a single protease is responsible for degradation of a particular antitoxin (although the proteases of interest comprise a family of related enzymes) (Van Melderen *et al.*, 1994; Lehnher & Yarmolinsky, 1995; Aizenman *et al.*, 1996; Christensen *et al.*, 2001; 2004; Kawano *et al.*, 2007; Christensen-Dalsgaard *et al.*, 2010; Donegan *et al.*, 2010). Degradation of the antitoxin component leads to subsequent toxin activation and increase in operon transcription in response to a toxin and antitoxin level imbalance. However, a halt of translation, induced for example by antibiotics, acts as another way of toxin activation by causing a drop in the production of labile antitoxin.

The significant influence of the TA systems on bacterial metabolism implies multiple ways of their activity



**Figure 2.** Binding of toxin-antitoxin complex to regulatory sequences leads to autorepression of TA operon expression

regulation. A well documented mechanism is the relation between the *mazEF* system of *E. coli* and locus *relA*, which codes for ATP:GTP 3'-diphosphotransferase implicated in the synthesis of 3',5'-guanosine bisphosphate (Justesen *et al.*, 1986; Metzger *et al.*, 1988). The ppGpp molecule is a signal of amino-acid starvation (Cashel, 1975; Gallant *et al.*, 1976). The *mazEF* locus is located downstream the *relA* locus (Masuda *et al.*, 1993) and is cotranscribed when *relA* expression is activated (Aizenman *et al.*, 1996; Christensen *et al.*, 2003; Hazan & Engelberg-Kulka, 2004). A similar neighbourhood pattern of the *mazEF* and *parDE* systems is found in genomes of other enteric bacteria such as *Shigella* and *Salmonella* (Pandey & Gerdes, 2005). Another example is the SOS system and its relations with various TA systems of *E. coli*. In this case the activation of SOS system leads to switching on the activity of TA systems including *hokE* (Fernandez De Henestrosa *et al.*, 2000), *yafNO* (McKenzie *et al.*, 2003; Christensen-Dalsgaard *et al.*, 2010), *tisAB* (Vogel *et al.*, 2004; Unoson & Wagner, 2008), *symER* (Kawano *et al.*, 2007), and *yefQ* (Motiejunaite *et al.*, 2007). A similar situation was recently reported for another *E. coli* TA system — *yafNO* (Singletary *et al.*, 2009).

The activity of TA systems can also be induced by systems responsible for *quorum sensing*. Such a mechanism has been reported for the *mazEF* system of *E. coli* (Kolodkin-Gal *et al.*, 2007). Another noteworthy fact is the possibility of cascade activation of TA systems (Hazan *et al.*, 2001) since the bacteria often carry more than a single TA system within their genome. Activation of a single system which leads to protein synthesis inhibition and subsequent activation of another TA system is plausible. An even more complex relation has been described for the *ygiUT* (*MqsR4*) system of *E. coli*. In this case activation of the TA system is necessary for activation of toxin CspD, whose gene promoter is controlled by the *ygiU/ygiT* (*MqsR/MqsA*) complex (Brown *et al.*, 2009; Kim *et al.*, 2010). Furthermore, a cross-regulation has been observed for homologous systems present in the genome (Yang *et al.*, 2010), where toxin-antitoxin complexes of one system bind to regulatory sequences of another TA system operon.

## FUNCTIONS OF CLASS II TA SYSTEMS

A plasmid maintenance function was initially assigned to several newly discovered plasmid-borne TA systems (Gerdes & Molin, 1986; Saurugger, 1986; Bravo *et al.*, 1988; Tsuchimoto *et al.*, 1988; Gerlitz *et al.*, 1990; Sobecky *et al.*, 1996). Cells that do not inherit a copy of a plasmid upon division do not survive the effect of a stable toxin after degradation of a labile antitoxin. Moreover, a role of multiple TA *loci* in stabilization of a megaintegron of *Vibrio cholerae* has been suggested (Pandey & Gerdes, 2005). There is no doubt that TA systems play a role in the phenomenon of mobile genetic element stabilization but operons of many TA systems are also located in the bacterial chromosome. Recent studies report that TA systems are mainly concerned with the regulation of bacterial metabolism rather than simple plasmid maintenance functions.

Toxin activity leads primarily to bacterial metabolic dormancy that can be abolished at initial stages (Nystrom, 1999; Pedersen *et al.*, 2002; Keren *et al.*, 2004; Gerdes *et al.*, 2005; Suzuki *et al.*, 2005; Buts *et al.*, 2005; Lewis, 2005; Inouye, 2006; Schumacher *et al.*, 2009; Fu *et al.*, 2009; Kasari *et al.*, 2010), which contrasts with ear-

lier suggestions that this activity leads to immediate cell death (Aizenman *et al.*, 1996; Hazan & Engelberg-Kulka, 2004; Engelberg-Kulka *et al.*, 2005). There are examples of such systems whose major role is to kill the cells, but this is only true in some specialized situations. A good example are formation of fruiting bodies of *Mycococcus xanthus* (Nariya & Inouye, 2008) or defence against phage infection in lactic acid bacteria (Forde & Fitzgerald, 1999). The question whether TA system activity leading to death of selected cells in a colony is a manifestation of an altruistic or other mechanism is currently a topic of discussion (Aizenman *et al.*, 1996; Forde & Fitzgerald, 1999; Nystrom, 1999; Lioy *et al.*, 2006).

A flexible response of a bacterial cell to stress conditions seems to be the major function of most TA systems. A reversible metabolic dormancy caused by their activation allows a bacterial cell to survive detrimental conditions. This phenomenon provides clear advantages in the case of starvation (Christensen *et al.*, 2001; Jorgensen *et al.*, 2009) as well as heat, osmotic and free-radicals-induced stress (Pedersen *et al.*, 2002; Senn *et al.*, 2005). Moreover, TA systems can contribute to the formation of persistent cells during an exposure to antibiotics (Falla & Chopra, 1998; Keren *et al.*, 2004; Dorr *et al.*, 2010; Kasari *et al.*, 2010). The mechanism of described phenomenon is straightforward in the case of drugs acting as transcription (eg. rifampicin) or translation (eg. chloramphenicol, doxycyclin, spectinomycin, erythromycin) inhibitors when the decay of the labile antitoxin causes the toxin activation. Paradoxically, antibiotics that are gyrase inhibitors (quinolone antibiotics) can act in a way similar to the *ccdAB* TA system, in which the toxin is a gyrase inhibitor. In this case binding of the inhibitor to an open gyrase–DNA complex induces DNA nicks (Drlica & Zhao, 1997; Jiang *et al.*, 2002), which is followed by SOS-system activation (Little & Mount, 1982; Karoui *et al.*, 1983; Bailone *et al.*, 1985). The same mechanism is proposed for homologues of *parDE* system (Hallez *et al.*, 2010). The described sequence of events leads to increased genetic diversity of a colony and may contribute to persisters formation (Couturier *et al.*, 1998) in the same way as do quinolone antibiotics (Drlica & Zhao, 1997).

The activity of TA systems can also modulate the behaviour of a bacterial colony. An increase in the expression of genes related to cell motility and structural genes of flagella has been reported for the *ygiUT* (*MqsRA*) system (Gonzalez Barrios, 2006). In turn the *hipAB* system is implicated in biofilm formation providing multi drug resistance (Lewis, 2007; 2008). TA systems can modulate formation of a biofilm over time (Kim *et al.*, 2009). In line with that, a recent report indicates elevated expression of TA systems in bacterial cells building a biofilm (Mitchell *et al.*, 2010).

A precise control over pathogenesis progression has been demonstrated for mRNA interferases exhibiting sequence specificity. This specificity allows for molecular evolution of target gene sequences. The mRNA interferases of the *mazEF-mt3* and *mazEF-mt7* systems are able to specifically recognize pentanucleotide sequences. In both cases a statistically significant representation of genes implicated in pathogenesis was found among genes containing underrepresented number of the recognized sequences (Zhu *et al.*, 2008). Such genes are resistant to the interferase activity and thereby are expected to be expressed even when the TA system is activated. A similar relation was found for the *sraP* gene of *S. aureus*. This gene, coding for a protein responsible for ad-

hesion to platelets (Siboo *et al.*, 2005), is characterized by a statistically significant overrepresentation of the sequence recognized by the mRNA interferase of the *mazEF<sub>Sa</sub>* TA system (Zhu *et al.*, 2009), hence its expression is suggested to be primarily turned off upon TA system activation. Additionally, the mentioned TA system may potentially be implicated in pathogenesis progression in yet another way. Downstream of the *mazEF<sub>Sa</sub>* locus a *sigB* locus is located (Kullik *et al.*, 1998; Gertz *et al.*, 1999; Ferreira *et al.*, 2004). The *sigB*-encoded alternative subunit  $\sigma^B$  of the RNA polymerase is responsible for global transcription regulation of virulence factors, comprising one of the most important staphylococcal systems of gene regulation responsible for pathogenesis (Wu *et al.*, 1996). In stress conditions the *sigB* locus is coexpressed with *mazEF<sub>Sa</sub>* (Senn *et al.*, 2005; Fu *et al.*, 2007; Donegan & Cheung, 2009). However, any potential functional relation demands further investigation since the elevated expression of *sigB* locus does not necessarily lead to a direct increase in the level of  $\sigma^B$  subunit (Senn *et al.*, 2005). Among other pathogenic strains also *Bacillus anthracis* possesses a TA system of the *mazEF* family, namely a *pemIK* module (Agarwal *et al.*, 2007; 2009). Recently a *pemIK* homologue located in a plasmid of an avian strains of *S. aureus* has been documented (Lowder *et al.*, 2009; Bukowski *et al.*, 2010). In this system the toxin is a sequence-specific endoribonuclease which targets a tetranucleotide sequence. Bioinformatic analysis of the occurrence of the recognized sequence in the coding sequences of the *S. aureus* genome elucidated a potential relation of the system with virulence factor regulation (Bukowski *et al.*, 2010).

## CLASS II TA SYSTEMS AS BIOTECHNOLOGICAL TOOLS

Two of the best-described TA systems have found application in molecular biology, namely *ccdAB* and *mazEF*. The former is used as a factor for positive selection of transformants, primarily in *E. coli* strains (Bernard *et al.*, 1994). Such systems, which are commercially available (e.g. StabyCloning™ and StabyExpress™, Delphi Genetics SA), are based on CcdB toxicity against gyrase and allow one-step selection of transformants ensuring stable vector plasmid maintenance (Fig. 3). This idea was originally developed by Szpirer and Milinkovitch (2005) followed by other efforts to develop a more complex system allowing increased production of recombinant protein (Stieber *et al.*, 2008).




Figure 3. *ccdAB* system components as tools for positive selection during cloning

The *mazEF* system has been adapted for single protein production (SPP) systems. The initial idea uses MazF toxin to trigger bacteriostasis and bacterial protein shutdown. The recombinant gene lacks the ACA sequences, recognized by the MazF interferase, therefore upon induction of MazF expression production of the recombinant protein of interest is continued almost exclusively. Moreover, bacteriostasis allows for culturing of the transformed strains in lower medium volumes than in traditional methods (Suzuki *et al.*, 2005; 2007). This idea has been successfully applied for protein production for NMR studies in 150-fold concentrated cultures, which allowed significant cost saving on isotopes (Mao *et al.*, 2009; Schneider *et al.*, 2009). Recently the SPP system based on MazF activity was extended with the capability for induction of protein production using particular amino acids. MazF mutants with histidine or tryptophan substitution were used in histidine or tryptophan auxotrophs, respectively. After transferring cells to the medium enriched in isotopes but lacking one of these amino acids the production of MazF is still provided. Subsequent addition of the amino acid induces exclusive production of the recombinant protein, since production of host proteins is blocked by the toxic action of MazF. Therefore, this approach allows not only single protein production but also high-efficiency isotope-labelling of the target protein (Vaiphei *et al.*, 2010).

TA systems are successfully used also in studies on eukaryotic cells. Recently a report concerning the usage of *mazEF* system in studies on HIV virus was published (Chono *et al.*, 2010). Further possible applications have already been suggested, such as TA-based contamination control in fermentation processes (Kristoffersen *et al.*, 2000), antibacterial drug development (Engelberg-Kulka *et al.*, 2004; Moritz & Hergenrother, 2007; Liroy *et al.*, 2010), selectable elimination of cells in cell cultures, tissue cultures and whole organisms (de la Cueva-Mendez *et al.*, 2003) or stable plasmid maintenance without antibiotic pressure (Wladyka *et al.*, 2010).

## CONCLUDING REMARKS

Results collected so far give a complex but concise image of the role of TA systems in bacterial physiology. Their functions range far beyond stabilization of mobile genetic elements. Metabolic dormancy induced by the systems seems a general but adequate response to various stress stimuli coming from the environment. Endoribonucleases, also termed mRNA interferases, are the most common group among the toxic components of various TA systems. Their activity leads to bacteriostasis through the inhibition of translation, which enables survival during starvation or antibiotic exposition. Further specialisation of interferases in selective sequence recognition allowed some genes to escape from expression suppression or, conversely, become exceptionally sensitive to a particular TA system. These phenomena are suggested to play a significant role in pathogen–host interaction and pathogenesis progression by modulation of biofilm formation and interactions with host proteins or coupling with other pathogen invasion-facilitating systems.

The relations among the ten families of class II TA systems are difficult to untangle. These TA systems are spread throughout the two huge domains of *Archaea* and *Bacteria*. Beside clear relationships, it seems that the similar way of acting and regulation of various groups of TA

systems are due to convergence. Components of such systems could have evolved divergently from unrelated groups of genes to create autoregulated operons coding for pairs of toxic protein and its inhibitor.

The physiological functions of the TA systems became a base for their successful applications as molecular biology tools, both in industry and research. Primarily they facilitate maintenance of plasmid vectors and transformant selection, but also effective overexpression of recombinant proteins. The potential application of TA systems in antibiotic therapy cannot be omitted as it is known that TA systems induce bacteriostasis, whose prolongation results in bacterial cell death. With the growing knowledge of TA systems new useful applications are expected to be developed.

## Acknowledgements

The authors thank Professor Adam Dubin for critical review of this manuscript.

This work was supported in part by grant NN302 130734 from the Ministry of Science and Higher Education.

## REFERENCES

- Agarwal S, Agarwal S, Bhatnagar R (2007) Identification and characterization of a novel toxin-antitoxin module from *Bacillus anthracis*. *FEBS Lett* **581**: 1727–1734.
- Agarwal S, Mishra NK, Bhatnagar S, Bhatnagar R (2009) PemK toxin of *Bacillus anthracis* is a ribonuclease: an insight into its active site, structure, and function. *J Biol Chem* **285**: 7254–7270.
- Aizenman E, Engelberg-Kulka H, Glaser G (1996) An *Escherichia coli* chromosomal “addiction module” regulated by guanosine 3',5'-bispyrophosphate: a model for programmed bacterial cell death. *Proc Natl Acad Sci USA* **93**: 6059–6063.
- Anantharaman V, Aravind L (2003) New connections in the prokaryotic toxin-antitoxin network: relationship with the eukaryotic non-sense-mediated RNA decay system. *Genome Biol* **4**: R81.
- Arbing MA, Handelman SK, Kuzin AP, Verdon G, Wang C, Su M, Rothenbacher FP, Abashidze M, Liu M, Hurley JM, Xiao R, Acton T, Inouye M, Montelione GT, Woychik NA, Hunt JF (2010) Crystal structures of Phd-Doc, HigA, and YeeU establish multiple evolutionary links between microbial growth-regulating toxin-antitoxin systems. *Structure* **18**: 996–1010.
- Arcus VL, Rainey PB, Turner SJ (2005) The PIN-domain toxin-antitoxin array in mycobacteria. *Trends Microbiol* **13**: 360–365.
- Bailey SE, Hayes F (2009) Influence of operator site geometry on transcriptional control by the YefM-YoeB toxin-antitoxin complex. *J Bacteriol* **191**: 762–772.
- Bailone A, Sommer S, Devoret R (1985) Mini-F plasmid-induced SOS signal in *Escherichia coli* is RecBC dependent. *Proc Natl Acad Sci USA* **82**: 5973–5977.
- Bech FW, Jorgensen ST, Diderichsen B, Karlstrom OH (1985) Sequence of the *relB* transcription unit from *Escherichia coli* and identification of the *relB* gene. *EMBO J* **4**: 1059–1066.
- Bernard P, Gabant P, Bahassi EM, Couturier M (1994) Positive-selection vectors using the F plasmid *cadB* killer gene. *Gene* **148**: 71–74.
- Black DS, Kelly AJ, Mardis MJ, Moyed HS (1991) Structure and organization of *hip*, an operon that affects lethality due to inhibition of peptidoglycan or DNA synthesis. *J Bacteriol* **173**: 5732–5739.
- Black DS, Irwin B, Moyed HS (1994) Autoregulation of *hip*, an operon that affects lethality due to inhibition of peptidoglycan or DNA synthesis. *J Bacteriol* **176**: 4081–4091.
- Blower TR, Fineran PC, Johnson MJ, Toth IK, Humphreys DP, Salmond GP (2009) Mutagenesis and functional characterization of the RNA and protein components of the *toxIN* abortive infection and toxin-antitoxin locus of *Erwinia*. *J Bacteriol* **191**: 6029–6039.
- Bravo A, de Torriontegui G, Diaz R (1987) Identification of components of a new stability system of plasmid R1, ParD, that is close to the origin of replication of this plasmid. *Mol Gen Genet* **210**: 101–110.
- Bravo A, Ortega S, de Torriontegui G, Diaz R (1988) Killing of *Escherichia coli* cells modulated by components of the stability system ParD of plasmid R1. *Mol Gen Genet* **215**: 146–151.
- Brown BL, Grigoriu S, Kim Y, Arruda JM, Davenport A, Wood TK, Peti W, Page R (2009) Three dimensional structure of the MqsR:MqsA complex: a novel TA pair comprised of a toxin ho-

- mologous to RelE and an antitoxin with unique properties. *PLoS Pathog* **5**: e1000706.
- Budde PP, Davis BM, Yuan J, Waldor MK (2007) Characterization of a higBA toxin-antitoxin locus in *Vibrio cholerae*. *J Bacteriol* **189**: 491–500.
- Bukowski M, Wladyka B, Lyzen R, Szalewska-Palasz A, Rojowska A, Dubin G, Dubin A (2010) A novel mRNA interferase encoded in toxin-antitoxin system of *Staphylococcus aureus* targeting tetranucleotide sequence. *Acta Biochim Pol* **57** (Suppl. 4): 9.
- Buts L, Lah J, Dao-Thi MH, Wyns L, Loris R (2005) Toxin-antitoxin modules as bacterial metabolic stress managers. *Trends Biochem Sci* **30**: 672–679.
- Camacho AG, Misselwitz R, Behlke J, Ayora S, Welfle K, Meinhart A, Lara B, Saenger W, Welfle H, Alonso JC (2002) *In vitro* and *in vivo* stability of the epsilon2zeta2 protein complex of the broad host-range *Streptococcus pyogenes* pSM19035 addiction system. *Biol Chem* **383**: 1701–1713.
- Cashel M (1975) Regulation of bacterial ppGpp and pppGpp. *Annu Rev Microbiol* **29**: 301–318.
- Chono H, Matsumoto K, Tsuda H, Saito N, Lee K, Kim S, Shibata H, Ageyama N, Terao K, Yasutomi Y, Mineno J, Kim S, Inouye M, Kato I (2010) Acquisition of HIV-1 resistance in T lymphocytes using an ACA-specific *E. coli* mRNA interferase. *Hum Gene Ther*.
- Christensen-Dalsgaard M, Gerdes K (2006) Two higBA loci in the *Vibrio cholerae* superintegron encode mRNA cleaving enzymes and can stabilize plasmids. *Mol Microbiol* **62**: 397–411.
- Christensen-Dalsgaard M, Gerdes K (2008) Translation affects YoeB and MazF messenger RNA interferase activities by different mechanisms. *Nucleic Acids Res* **36**: 6472–6481.
- Christensen-Dalsgaard M, Jorgensen MG, Gerdes K (2010) Three new RelE-homologous mRNA interferases of *Escherichia coli* differentially induced by environmental stresses. *Mol Microbiol* **75**: 333–348.
- Christensen SK, Mikkelsen M, Pedersen K, Gerdes K (2001) RelE, a global inhibitor of translation, is activated during nutritional stress. *Proc Natl Acad Sci USA* **98**: 14328–14333.
- Christensen SK, Pedersen K, Hansen FG, Gerdes K (2003) Toxin-antitoxin loci as stress-response-elements: ChpAK/MazF and ChpBK cleave translated RNAs and are counteracted by tmRNA. *J Mol Biol* **332**: 809–819.
- Christensen SK, Maenhaut-Michel G, Mine N, Gottesman S, Gerdes K, Van Melderen L (2004) Overproduction of the Lon protease triggers inhibition of translation in *Escherichia coli*: involvement of the yefM-yoeB toxin-antitoxin system. *Mol Microbiol* **51**: 1705–1717.
- Correia FF, D'Onofrio A, Rejtar T, Li L, Karger BL, Makarova K, Koonin EV, Lewis K (2006) Kinase activity of overexpressed HipA is required for growth arrest and multidrug tolerance in *Escherichia coli*. *J Bacteriol* **188**: 8360–8367.
- Couturier M, Bahassi el M, Van Melderen L (1998) Bacterial death by DNA gyrase poisoning. *Trends Microbiol* **6**: 269–275.
- Daines DA, Wu MH, Yuan SY (2007) VapC-1 of nontypeable *Haemophilus influenzae* is a ribonuclease. *J Bacteriol* **189**: 5041–5048.
- de la Cueva-Mendez G, Mills AD, Clay-Farrace L, Diaz-Orejas R, Laskey RA (2003) Regulatable killing of eukaryotic cells by the prokaryotic proteins Kid and Kis. *EMBO J* **22**: 246–251.
- de la Hoz AB, Ayora S, Sitkiewicz I, Fernandez S, Pankiewicz R, Alonso JC, Ceglowski P (2000) Plasmid copy-number control and better-than-random segregation genes of pSM19035 share a common regulator. *Proc Natl Acad Sci USA* **97**: 728–733.
- DeLano WL (2002) The PyMOL Molecular Graphics System. *DeLano Scientific, San Carlos*.
- Diago-Navarro E, Hernandez-Arriaga AM, Lopez-Villarejo J, Munoz-Gomez AJ, Kamphuis MB, Boelens R, Lemonnier M, Diaz-Orejas R (2010) parD toxin-antitoxin system of plasmid R1 — basic contributions, biotechnological applications and relationships with closely-related toxin-antitoxin systems. *FEBS J* **277**: 3097–3117.
- Diderichsen B, Fiil NP, Lavallo R (1977) Genetics of the relB locus in *Escherichia coli*. *J Bacteriol* **131**: 30–33.
- Donegan NP, Cheung AL (2009) Regulation of the mazEF toxin-antitoxin module in *Staphylococcus aureus* and its impact on sigB expression. *J Bacteriol* **191**: 2795–2805.
- Donegan NP, Thompson ET, Fu Z, Cheung AL (2010) Proteolytic regulation of toxin-antitoxin systems by ClpPC in *Staphylococcus aureus*. *J Bacteriol* **192**: 1416–1422.
- Dorr T, Vulic M, Lewis K (2010) Ciprofloxacin causes persister formation by inducing the TisB toxin in *Escherichia coli*. *PLoS Biol* **8**: e1000317.
- Drlica K, Zhao X (1997) DNA gyrase, topoisomerase IV, and the 4-quinolones. *Microbiol Mol Biol Rev* **61**: 377–392.
- Engelberg-Kulka H, Sat B, Reches M, Amitai S, Hazan R (2004) Bacterial programmed cell death systems as targets for antibiotics. *Trends Microbiol* **12**: 66–71.
- Engelberg-Kulka H, Hazan R, Amitai S (2005) mazEF: a chromosomal toxin-antitoxin module that triggers programmed cell death in bacteria. *J Cell Sci* **118**: 4327–4332.
- Falla TJ, Chopra I (1998) Joint tolerance to beta-lactam and fluoroquinolone antibiotics in *Escherichia coli* results from overexpression of hipA. *Antimicrob Agents Chemother* **42**: 3282–3284.
- Fernandez De Henestrosa AR, Ogi T, Aoyagi S, Chafin D, Hayes JJ, Ohmori H, Woodgate R (2000) Identification of additional genes belonging to the LexA regulon in *Escherichia coli*. *Mol Microbiol* **35**: 1560–1572.
- Ferreira A, Gray M, Wiedmann M, Boor KJ (2004) Comparative genomic analysis of the sigB operon in *Listeria monocytogenes* and in other Gram-positive bacteria. *Curr Microbiol* **48**: 39–46.
- Fineran PC, Blower TR, Foulds IJ, Humphreys DP, Lilley KS, Salmon GP (2009) The phage abortive infection system, ToxIN, functions as a protein-RNA toxin-antitoxin pair. *Proc Natl Acad Sci USA* **106**: 894–899.
- Forde A, Fitzgerald GF (1999) Bacteriophage defence systems in lactic acid bacteria. *Antonie Van Leeuwenhoek* **76**: 89–113.
- Fu Z, Donegan NP, Memmi G, Cheung AL (2007) Characterization of MazF<sub>Sa</sub>, an endoribonuclease from *Staphylococcus aureus*. *J Bacteriol* **189**: 8871–8879.
- Fu Z, Tamber S, Memmi G, Donegan NP, Cheung AL (2009) Overexpression of MazF<sub>Sa</sub> in *Staphylococcus aureus* induces bacteriostasis by selectively targeting mRNAs for cleavage. *J Bacteriol* **191**: 2051–2059.
- Gallant J, Shell L, Bittner R (1976) A novel nucleotide implicated in the response of *E. coli* to energy source downshift. *Cell* **7**: 75–84.
- Galvani C, Terry J, Ishiguro EE (2001) Purification of the RelB and RelE proteins of *Escherichia coli*: RelE binds to RelB and to ribosomes. *J Bacteriol* **183**: 2700–2703.
- Gazit E, Sauer RT (1999) The Doc toxin and Phd antidote proteins of the bacteriophage P1 plasmid addiction system form a heterotrimeric complex. *J Biol Chem* **274**: 16813–16818.
- Gerdes K (2000) Toxin-antitoxin modules may regulate synthesis of macromolecules during nutritional stress. *J Bacteriol* **182**: 561–572.
- Gerdes K, Molin S (1986) Partitioning of plasmid R1. Structural and functional analysis of the parA locus. *J Mol Biol* **190**: 269–279.
- Gerdes K, Wagner EG (2007) RNA antitoxins. *Curr Opin Microbiol* **10**: 117–124.
- Gerdes K, Bech FW, Jorgensen ST, Lobner-Olesen A, Rasmussen PB, Atlung T, Boe L, Karlstrom O, Molin S, von Meyenburg K (1986) Mechanism of postsegregational killing by the hok gene product of the parB system of plasmid R1 and its homology with the relF gene product of the *E. coli* relB operon. *Embo J* **5**: 2023–2029.
- Gerdes K, Christensen SK, Lobner-Olesen A (2005) Prokaryotic toxin-antitoxin stress response loci. *Nat Rev Microbiol* **3**: 371–382.
- Gerlitz M, Hrabak O, Schwab H (1990) Partitioning of broad-host-range plasmid RP4 is a complex system involving site-specific recombination. *J Bacteriol* **172**: 6194–6203.
- Gertz S, Engelmann S, Schmid R, Ohlsen K, Hacker J, Hecker M (1999) Regulation of sigmaB-dependent transcription of sigB and asp23 in two different *Staphylococcus aureus* strains. *Mol Gen Genet* **261**: 558–566.
- Gonzalez Barrios AF, Zuo R, Hashimoto Y, Yang L, Bentley WE, Wood TK (2006) Autoinducer 2 controls biofilm formation in *Escherichia coli* through a novel motility quorum-sensing regulator (MqsR, B3022). *J Bacteriol* **188**: 305–316.
- Gottesman S (1996) Proteases and their targets in *Escherichia coli*. *Annu Rev Genet* **30**: 465–506.
- Greenfield TJ, Ehli E, Kirshenmann T, Franch T, Gerdes K, Weaver KE (2000) The antisense RNA of the par locus of pAD1 regulates the expression of a 33-amino-acid toxic peptide by an unusual mechanism. *Mol Microbiol* **37**: 652–660.
- Hallez R, Geeraerts D, Sterckx Y, Mine N, Loris R, Van Melderen L (2010) New toxins homologous to ParE belonging to three-component toxin-antitoxin systems in *Escherichia coli* O157:H7. *Mol Microbiol* **76**: 719–732.
- Hayes CS, Sauer RT (2003) Toxin-antitoxin pairs in bacteria: killers or stress regulators? *Cell* **112**: 2–4.
- Hazan R, Engelberg-Kulka H (2004) *Escherichia coli* mazEF-mediated cell death as a defense mechanism that inhibits the spread of phage P1. *Mol Genet Genomics* **272**: 227–234.
- Hazan R, Sat B, Reches M, Engelberg-Kulka H (2001) Postsegregational killing mediated by the P1 phage “addiction module” phd-doc requires the *Escherichia coli* programmed cell death system mazEF. *J Bacteriol* **183**: 2046–2050.
- Inouye M (2006) The discovery of mRNA interferases: implication in bacterial physiology and application to biotechnology. *J Cell Physiol* **209**: 670–676.
- Jensen SO, Apisiridej S, Kwong SM, Yang YH, Skurray RA, Firth N (2010) Analysis of the prototypical *Staphylococcus aureus* multiresistance plasmid pSK1. *Plasmid* **64**: 135–142.
- Jiang Y, Pogliano J, Helinski DR, Konieczny I (2002) ParE toxin encoded by the broad-host-range plasmid RK2 is an inhibitor of *Escherichia coli* gyrase. *Mol Microbiol* **44**: 971–979.
- Jorgensen MG, Pandey DP, Jaskolska M, Gerdes K (2009) HicA of *Escherichia coli* defines a novel family of translation-independent mRNA interferases in bacteria and archaea. *J Bacteriol* **191**: 1191–1199.

- Justesen J, Lund T, Skou Pedersen F, Kjeldgaard NO (1986) The physiology of stringent factor (ATP:GTP 3'-diphosphotransferase) in *Escherichia coli*. *Biochimie* **68**: 715–722.
- Kamada K, Hanaoka F, Burley SK (2003) Crystal structure of the MazE/MazF complex: molecular bases of antidote-toxin recognition. *Mol Cell* **11**: 875–884.
- Karoui H, Bex F, Dreze P, Couturier M (1983) Ham22, a mini-F mutation which is lethal to host cell and promotes recA-dependent induction of lambda-doid prophage. *EMBO J* **2**: 1863–1868.
- Kasari V, Kurg K, Margus T, Tenson T, Kaldalu N (2010) The *Escherichia coli* mqsR and ygiT genes encode a new toxin-antitoxin pair. *J Bacteriol* **192**: 2908–2919.
- Kawano M, Aravind L, Storz G (2007) An antisense RNA controls synthesis of an SOS-induced toxin evolved from an antitoxin. *Mol Microbiol* **64**: 738–754.
- Kedzierska B, Lian LY, Hayes F (2007) Toxin-antitoxin regulation: bimodal interaction of YefM-YoeB with paired DNA palindromes exerts transcriptional autorepression. *Nucleic Acids Res* **35**: 325–339.
- Keren I, Kaldalu N, Spoering A, Wang Y, Lewis K (2004) Persister cells and tolerance to antimicrobials. *FEMS Microbiol Lett* **230**: 13–8.
- Kim Y, Wang X, Ma Q, Zhang XS, Wood TK (2009) Toxin-antitoxin systems in *Escherichia coli* influence biofilm formation through YjgK (TabA) and fimbriae. *J Bacteriol* **191**: 1258–1267.
- Kim Y, Wang X, Zhang XS, Grigoriu S, Page R, Peti W, Wood TK (2010) *Escherichia coli* toxin/antitoxin pair MqsR/MqsA regulate toxin CspD. *Environ Microbiol* **12**: 1105–1121.
- Kolodkin-Gal I, Hazan R, Gaathon A, Carmeli S, Engelberg-Kulka H (2007) A linear pentapeptide is a quorum-sensing factor required for mazEF-mediated cell death in *Escherichia coli*. *Science* **318**: 652–655.
- Korch SB, Hill TM (2006) Ectopic overexpression of wild-type and mutant hipA genes in *Escherichia coli*: effects on macromolecular synthesis and persister formation. *J Bacteriol* **188**: 3826–3836.
- Kristoffersen P, Jensen GB, Gerdes K, Piskur J (2000) Bacterial toxin-antitoxin gene system as containment control in yeast cells. *Appl Environ Microbiol* **66**: 5524–5526.
- Kullik I, Giachino P, Fuchs T (1998) Deletion of the alternative sigma factor sigmaB in *Staphylococcus aureus* reveals its function as a global regulator of virulence genes. *J Bacteriol* **180**: 4814–4820.
- Lavalle R (1965) New mutants for regulation of RNA synthesis. *Bull Soc Chim Biol (Paris)* **47**: 1567–1570.
- Lehnherr H, Yarmolinsky MB (1995) Addiction protein Phd of plasmid prophage P1 is a substrate of the ClpXP serine protease of *Escherichia coli*. *Proc Natl Acad Sci USA* **92**: 3274–3277.
- Lehnherr H, Maguin E, Jafri S, Yarmolinsky MB (1993) Plasmid addiction genes of bacteriophage P1: doc, which causes cell death on curing of prophage, and phd, which prevents host death when prophage is retained. *J Mol Biol* **233**: 414–428.
- Lewis K (2005) Persister cells and the riddle of biofilm survival. *Biochemistry (Mosc)* **70**: 267–274.
- Lewis K (2007) Persister cells, dormancy and infectious disease. *Nat Rev Microbiol* **5**: 48–56.
- Lewis K (2008) Multidrug tolerance of biofilms and persister cells. *Curr Top Microbiol Immunol* **322**: 107–131.
- Li GY, Zhang Y, Inouye M, Ikura M (2008) Structural mechanism of transcriptional autorepression of the *Escherichia coli* RelB/RelE antitoxin/toxin module. *J Mol Biol* **380**: 107–119.
- Lioy VS, Martin MT, Camacho AG, Lurz R, Antelmann H, Hecker M, Hitchin E, Ridge Y, Wells JM, Alonso JC (2006) pSM19035-encoded zeta toxin induces stasis followed by death in a subpopulation of cells. *Microbiology* **152**: 2365–2379.
- Lioy VS, Rey O, Balsa D, Pellicer T, Alonso JC (2010) A toxin-antitoxin module as a target for antimicrobial development. *Plasmid* **63**: 31–39.
- Little JW, Mount DW (1982) The SOS regulatory system of *Escherichia coli*. *Cell* **29**: 11–22.
- Liu M, Zhang Y, Inouye M, Woychik NA (2008) Bacterial addiction module toxin Doc inhibits translation elongation through its association with the 30S ribosomal subunit. *Proc Natl Acad Sci USA* **105**: 5885–5890.
- Lowder BV, Guinane CM, Ben Zakour NL, Weinert LA, Conway-Morris A, Cartwright RA, Simpson AJ, Rambaut A, Nubel U, Fitzgerald JR (2009) Recent human-to-poultry host jump, adaptation, and pandemic spread of *Staphylococcus aureus*. *Proc Natl Acad Sci USA* **106**: 19545–19550.
- Magnuson R, Yarmolinsky MB (1998) Corepression of the P1 addiction operon by Phd and Doc. *J Bacteriol* **180**: 6342–6351.
- Magnuson R, Lehnherr H, Mukhopadhyay G, Yarmolinsky MB (1996) Autoregulation of the plasmid addiction operon of bacteriophage P1. *J Biol Chem* **271**: 18705–18710.
- Makarova KS, Grishin NV, Koonin EV (2006) The HicAB cassette, a putative novel, RNA-targeting toxin-antitoxin system in archaea and bacteria. *Bioinformatics* **22**: 2581–2584.
- Makarova KS, Wolf YI, Koonin EV (2009) Comprehensive comparative-genomic analysis of type 2 toxin-antitoxin systems and related mobile stress response systems in prokaryotes. *Biol Direct* **4**: 19.
- Mao L, Tang Y, Vaiphei ST, Shimazu T, Kim SG, Mani R, Fakhoury E, White E, Montelione GT, Inouye M (2009) Production of membrane proteins for NMR studies using the condensed single protein (cSPP) production system. *J Struct Funct Genomics* **10**: 281–289.
- Marianovsky I, Aizenman E, Engelberg-Kulka H, Glaser G (2001) The regulation of the *Escherichia coli* mazEF promoter involves an unusual alternating palindrome. *J Biol Chem* **276**: 5975–5984.
- Masuda Y, Miyakawa K, Nishimura Y, Ohtsubo E (1993) chpA and chpB, *Escherichia coli* chromosomal homologs of the pem locus responsible for stable maintenance of plasmid R100. *J Bacteriol* **175**: 6850–6856.
- McKenzie GJ, Magner DB, Lee PL, Rosenberg SM (2003) The dinB operon and spontaneous mutation in *Escherichia coli*. *J Bacteriol* **185**: 3972–3977.
- Meinhart A, Alonso JC, Strater N, Saenger W (2003) Crystal structure of the plasmid maintenance system epsilon/zeta: functional mechanism of toxin zeta and inactivation by epsilon 2 zeta 2 complex formation. *Proc Natl Acad Sci USA* **100**: 1661–1666.
- Metzger S, Dror IB, Aizenman E, Schreiber G, Toone M, Friesen JD, Cashel M, Glaser G (1988) The nucleotide sequence and characterization of the *relA* gene of *Escherichia coli*. *J Biol Chem* **263**: 15699–15704.
- Miki T, Park JA, Nagao K, Murayama N, Horiuchi T (1992) Control of segregation of chromosomal DNA by sex factor F in *Escherichia coli*. Mutants of DNA gyrase subunit A suppress letD (ccdB) product growth inhibition. *J Mol Biol* **225**: 39–52.
- Miller S (1989) The structure of interfaces between subunits of dimeric and tetrameric proteins. *Protein Eng* **3**: 77–83.
- Mitchell HL, Dashper SG, Catmull DV, Paolini RA, Cleal SM, Slakeski N, Tan KH, Reynolds EC (2010) *Treponema denticola* biofilm-induced expression of a bacteriophage, toxin-antitoxin systems and transposases. *Microbiology* **156**: 774–788.
- Mittenhuber G (1999) Occurrence of mazEF-like antitoxin/toxin systems in bacteria. *J Mol Microbiol Biotechnol* **1**: 295–302.
- Moritz EM, Hergenrother PJ (2007) Toxin-antitoxin systems are ubiquitous and plasmid-encoded in vancomycin-resistant enterococci. *Proc Natl Acad Sci USA* **104**: 311–316.
- Mosteller RD (1978) Evidence that glucose starvation-sensitive mutants are altered in the *relB* locus. *J Bacteriol* **133**: 1034–1037.
- Motiejunaite R, Armalyte J, Markuckas A, Suziedeliene E (2007) *Escherichia coli* dinJ-yafQ genes act as a toxin-antitoxin module. *FEMS Microbiol Lett* **268**: 112–119.
- Munoz-Gomez AJ S-SS, Berzal-Herranz A, Lemonnier M, Diaz RO (2004) Insights into the specificity of RNA cleavage by the *Escherichia coli* MazF toxin. *FEBS Lett* **567**: 316–320.
- Nariya H, Inouye M (2008) MazF, an mRNA interferase, mediates programmed cell death during multicellular *Mycobacterium* development. *Cell* **132**: 55–66.
- Nystrom T (1999) Starvation, cessation of growth and bacterial aging. *Curr Opin Microbiol* **2**: 214–219.
- Ogura T, Hiraga S (1983) Mini-F plasmid genes that couple host cell division to plasmid proliferation. *Proc Natl Acad Sci USA* **80**: 4784–4788.
- Pandey DP, Gerdes K (2005) Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. *Nucleic Acids Res* **33**: 966–976.
- Pedersen K, Christensen SK, Gerdes K (2002) Rapid induction and reversal of a bacteriostatic condition by controlled expression of toxins and antitoxins. *Mol Microbiol* **45**: 501–510.
- Pedersen K, Zavialov AV, Pavlov MY, Elf J, Gerdes K, Ehrenberg M (2003) The bacterial toxin RelE displays codon-specific cleavage of mRNAs in the ribosomal A site. *Cell* **112**: 131–140.
- Robson J, McKenzie JL, Cursons R, Cook GM, Arcus VL (2009) The vapBC operon from *Mycobacterium smegmatis* is an autoregulated toxin-antitoxin module that controls growth via inhibition of translation. *J Mol Biol* **390**: 353–367.
- Saurugger PN, Hrabak O, Schwab H, Lafferty RM (1986) Mapping and cloning of the par-region of broad-host-range plasmid RP4. *J Biotechnol* **4**: 333–343.
- Schneider WM, Inouye M, Montelione GT, Roth MJ (2009) Independently inducible system of gene expression for condensed single protein production (cSPP) suitable for high efficiency isotope enrichment. *J Struct Funct Genomics* **10**: 219–225.
- Schumacher MA, Piro KM, Xu W, Hansen S, Lewis K, Brennan RG (2009) Molecular mechanisms of HipA-mediated multidrug tolerance and its neutralization by HipB. *Science* **323**: 396–401.
- Senn MM, Giachino P, Homerova D, Steinhuber A, Strassner J, Kormanec J, Fluckiger U, Berger-Bachi B, Bischoff M (2005) Molecular analysis and organization of the sigmaB operon in *Staphylococcus aureus*. *J Bacteriol* **187**: 8006–8019.
- Sevin EW, Barloy-Hubler F (2007) RASTA-Bacteria: a web-based tool for identifying toxin-antitoxin loci in prokaryotes. *Genome Biol* **8**: R155.
- Siboo IR, Chambers HF, Sullam PM (2005) Role of SraP, a serine-rich surface protein of *Staphylococcus aureus*, in binding to human platelets. *Infect Immun* **73**: 2273–2280.



- Singletary LA, Gibson JL, Tanner EJ, McKenzie GJ, Lee PL, Gonzalez C, Rosenberg SM (2009) An SOS-regulated type 2 toxin-antitoxin system. *J Bacteriol* **191**: 7456–7465.
- Sobecky PA, Easter CL, Bear PD, Helinski DR (1996) Characterization of the stable maintenance properties of the par region of broad-host-range plasmid RK2. *J Bacteriol* **178**: 2086–2093.
- Stieber D, Gabant P, Szpirer C (2008) The art of selective killing: plasmid toxin/antitoxin systems and their technological applications. *Biotechniques* **45**: 344–346.
- Suzuki M, Zhang J, Liu M, Woychik NA, Inouye M (2005) Single protein production in living cells facilitated by an mRNA interferase. *Mol Cell* **18**: 253–261.
- Suzuki M, Mao L, Inouye M (2007) Single protein production (SPP) system in *Escherichia coli*. *Nat Protoc* **2**: 1802–1810.
- Szpirer CY, Milinkovitch MC (2005) Separate-component-stabilization system for protein and DNA production without the use of antibiotics. *Biotechniques* **38**: 775–781.
- Thisted T, Gerdes K (1992) Mechanism of post-segregational killing by the *hok/sok* system of plasmid R1. *Sok* antisense RNA regulates *hok* gene expression indirectly through the overlapping *mok* gene. *J Mol Biol* **223**: 41–54.
- Tsilibaris V, Maenhaut-Michel G, Mine N, Van Melderen L (2007) What is the benefit to *Escherichia coli* of having multiple toxin-antitoxin systems in its genome? *J Bacteriol* **189**: 6101–6108.
- Tsuchimoto S, Ohtsubo E (1993) Autoregulation by cooperative binding of the PemI and PemK proteins to the promoter region of the pem operon. *Mol Gen Genet* **237**: 81–88.
- Tsuchimoto S, Ohtsubo H, Ohtsubo E (1988) Two genes, pemK and pemI, responsible for stable maintenance of resistance plasmid R100. *J Bacteriol* **170**: 1461–1466.
- Unoson C, Wagner EG (2008) A small SOS-induced toxin is targeted against the inner membrane in *Escherichia coli*. *Mol Microbiol* **70**: 258–270.
- Vaiphei ST, Mao L, Shimazu T, Park JH, Inouye M (2010) Use of amino acids as inducers for high-level protein expression in the single-protein production system. *Appl Environ Microbiol* **76**: 6063–6068.
- Van Melderen L, Bernard P, Couturier M (1994) Lon-dependent proteolysis of CcdA is the key control for activation of CcdB in plasmid-free segregant bacteria. *Mol Microbiol* **11**: 1151–1157.
- Van Melderen L, Saavedra De Bast M (2009) Bacterial toxin-antitoxin systems: more than selfish entities? *PLoS Genet* **5**: e1000437.
- Vogel J, Argaman L, Wagner EG, Altuvia S (2004) The small RNA IstR inhibits synthesis of an SOS-induced toxic peptide. *Curr Biol* **14**: 2271–2276.
- Weaver KE, Reddy SG, Brinkman CL, Patel S, Bayles KW, Endres JL (2009) Identification and characterization of a family of toxin-antitoxin systems related to the *Enterococcus faecalis* plasmid pAD1 par addiction module. *Microbiology* **155**: 2930–2940.
- Władyka B, Ilczyszyn WM, Pogwizd J, Rojowska A, Malachowa J, Bonar E, Polakowska K, Dubin G, Dubin A (2010) Potential application of staphylococcal pCH91 plasmid in biotechnology. *Acta Biochim Pol* **57** (Suppl. 4): 24.
- Wu S, de Lencastre H, Tomasz A (1996) Sigma-B, a putative operon encoding alternate sigma factor of *Staphylococcus aureus* RNA polymerase: molecular cloning and DNA sequencing. *J Bacteriol* **178**: 6036–6042.
- Yamaguchi Y, Park JH, Inouye M (2009) MqsR, a crucial regulator for quorum sensing and biofilm formation, is a GCU-specific mRNA interferase in *Escherichia coli*. *J Biol Chem* **284**: 28746–28753.
- Yang M, Gao C, Wang Y, Zhang H, He ZG (2010) Characterization of the interaction and cross-regulation of three *Mycobacterium tuberculosis* RelBE modules. *PLoS One* **5**: e10672.
- Zhang Y, Inouye M (2009) The inhibitory mechanism of protein synthesis by YoeB, an *Escherichia coli* toxin. *J Biol Chem* **284**: 6627–6638.
- Zhang J, Zhang Y, Zhu L, Suzuki M, Inouye M (2004) Interference of mRNA function by sequence-specific endoribonuclease PemK. *J Biol Chem* **279**: 20678–20684.
- Zhang Y, Zhang J, Hoefflich KP, Ikura M, Qing G, Inouye M (2003) MazF cleaves cellular mRNAs specifically at ACA to block protein synthesis in *Escherichia coli*. *Mol Cell* **12**: 913–923.
- Zhang Y, Zhu L, Zhang J, Inouye M (2005) Characterization of ChpBK, an mRNA interferase from *Escherichia coli*. *J Biol Chem* **280**: 26080–26088.
- Zhu L, Zhang Y, Teh JS, Zhang J, Connell N, Rubin H, Inouye M (2006) Characterization of mRNA interferases from *Mycobacterium tuberculosis*. *J Biol Chem* **281**: 18638–18643.
- Zhu L, Phadtare S, Nariya H, Ouyang M, Husson RN, Inouye M (2008) The mRNA interferases, MazF-mt3 and MazF-mt7 from *Mycobacterium tuberculosis* target unique pentad sequences in single-stranded RNA. *Mol Microbiol* **69**: 559–569.
- Zhu L, Inoue K, Yoshizumi S, Kobayashi H, Zhang Y, Ouyang M, Kato F, Sugai M, Inouye M (2009) *Staphylococcus aureus* MazF specifically cleaves a pentad sequence, UACAU, which is unusually abundant in the mRNA for pathogenic adhesive factor SraP. *J Bacteriol* **191**: 3248–3255.