

## Interaction of HIV Tat model peptides with tRNA and 5S rRNA

Małgorzata Giel-Pietraszuk<sup>1</sup>, Mirosława Z. Barciszewska<sup>1</sup>, Piotr Mucha<sup>2</sup>, Piotr Rekowski<sup>2</sup>, Gotfryd Kupryszewski<sup>2</sup> and Jan Barciszewski<sup>1</sup>

<sup>1</sup>*Institute of Bioorganic Chemistry, Polish Academy of Sciences, Z. Noskowskiego 12/14, 61-704 Poznań, Poland;*

<sup>2</sup>*Faculty of Chemistry, University of Gdańsk, J. Sobieskiego 18, 80-952 Gdańsk, Poland*

**Key words:** Tat-peptides, TAR RNA, ribosomal 5S RNA, tRNA<sup>Phe</sup>, protein-nucleic acid interactions

New data are presented on the interaction of model synthetic peptides containing an arginine-rich region of human immunodeficiency virus (HIV-Tat), with native RNA molecules: tRNA<sup>Phe</sup> of *Saccharomyces cerevisiae* and 5S rRNA from *Lupinus luteus*. Both RNA species form complexes with the Tat1 (GRKKRRQRRRA) and Tat2 (GRKKRRQRRRAPQDSQTHQASLSKQPA) peptides, as shown by electrophoretic gel shift and RNase footprint assays, and CD measurements. The nucleotide sequence UGGG located in the dihydrouridine loop of tRNA<sup>Phe</sup> as well as in the loop D of 5S rRNA is specifically protected against RNases. Our data indicate direct interactions of guanine of RNA moieties with arginine residues. These interactions seem similar to those observed in DNA-protein complexes, but different from those previously observed in the TAR RNA-Tat complexes.

Over recent years the mechanism of protein-nucleic acid interactions in solutions and crystal state was the subject of numerous studies. On the basis of crystallographic data several protein structural motifs responsible for specific recognition of DNA have been found [1]. A recurring feature in all these domains is a recognitory  $\alpha$ -helix, which binds to, and perfectly fits into, the major groove of B-DNA where amino-acid side chains form specific hydrogen bonds and non-specific contacts with bases and backbone phosphates, respectively. In some cases, high specificity of binding of DNA to protein through bound water molecules has been achieved [2]. Detailed analysis of crystal structures of various complexes has allowed several groups to formulate a "code" of interaction of protein transcription factors with DNA [3-5]. Gener-

ally, it predicts that the guanine and adenine residues form specific contacts with arginine and glutamine moiety, respectively. However, the data collected up to now do not confirm the existence of a similar coding system for RNA-protein complexes [6].

Recent advances in understanding of RNA-protein interactions come mostly from studies of three dimensional structure of the complexes of glutaminyl-, aspartyl- and seryl-tRNA synthetases with their cognate tRNAs [7-9], R17 coat protein with a 19 nucleotide long RNA hairpin [10] and U1A ribonucleo-protein domain with a 20 nucleotide long RNA hairpin [11]. A general conclusion drawn from those data indicates that it is rather the tertiary structure than nucleotide sequence of RNA which is the most important feature for its specific recognition by a cog-

\*This study was supported by a grant from the State Committee for Scientific Research (KBN) No. PB 218/T09/95/09.

**Abbreviations:** ARM, arginine-rich motif; BIV, bovine immunodeficiency virus; FZ-CE, free zone capillary electrophoresis; TAR RNA, transactivation response RNA.

nate protein. On the protein part, an RNA-protein binding motif is responsible for specific interaction with ribonucleic acid. One of such domains consists of a short string of basic amino acids which contains mainly arginine residues (arginine-rich motif, ARM) [6, 12]. Similar amino-acid stretches have been identified also in other proteins, e.g. as bacterial antiterminators, ribosomal proteins, coat proteins of RNA viruses, human immunodeficiency virus Tat (HIV-Tat) and Rev (HIV-Rev) proteins [13], and a bovine immunodeficiency virus Tat (BIV-Tat) protein [14]. A Tat protein activates expression of the HIV-1 genome by its specific binding to the transactivation response RNA (TAR RNA) sequence, located at the 5' end of the untranslated leader region of HIV mRNA [15]. The latter acquires a specific secondary structure involving a helical stem interrupted by a trinucleotide bulge ended with a six-residue loop [15]. Some time ago the complex structure of TAR RNA complexed with arginine amide was solved by NMR spectroscopy [16]. Hydrogen bonds between arginine and guanine residues in position 26 and two phosphates (positions 22 and 23) have been identified. The binding of arginine induces structural changes in RNA with formation of a U23-A27-U38 base triplet [16, 17]. These data have not been confirmed in other studies of the HIV RNA-Tat, where the base triplet has not been observed [18]. The computer modeling studies of TAR RNA-Tat complex have demonstrated that it is possible to accommodate the  $\alpha$ -helix of peptide into the RNA major groove, widened significantly by the bulge [19]. In that location of the  $\alpha$ -helix of Tat-peptide, the guanidine group of arginine fits to the binding pocket, and side chains of amino acid interact with A22 phosphoryl group and O6 oxygen of G26, but have hydrogen bonds with the N7 atom of A22 and the O4 atom of U23. The proposed interactions in this model differ slightly from the model proposed by Puglisi *et al.* [16] where guanidine group of arginine forms hydrogen bonds with N7 and O6 atoms of G26, and electrostatic interactions with U23 and A22 phosphoryl group.

In order to determine which RNA domain binds to arginine rich peptides, we used yeast

tRNA<sup>Phe</sup> and 5S rRNA from yellow lupin as model RNA molecules. These RNA molecules were chosen because they have a 5'-UGGG sequence in some of their loops identical to that in TAR RNA. Moreover, the three dimensional structure of yeast tRNA<sup>Phe</sup> is well known for over 20 years [20] and the tertiary structure of 5S rRNA has also been proposed [21]. As model proteins we used two short synthetic peptides with amino-acid sequence corresponding to the arginine-rich RNA-binding domain of HIV-1 Tat. In this paper we have shown by electrophoretic, CD and RNases protection assays that RNA molecules form complexes with these peptides, the Tat-peptides bind to the dihydrouridine loop of tRNA<sup>Phe</sup> and to the D-loop of 5S rRNA. These interactions resemble very much the guanine-arginine recognition mode observed previously in numerous DNA-protein complexes. The data presented here are at variance with those proposed earlier for TAR-Tat, in which protein was suggested to recognize a distorted helical RNA fragment [15].

## EXPERIMENTAL

### Synthesis of polypeptides

Tat1 and Tat2 peptides, having the amino-acid sequences shown in Table 1, were synthesized manually on a cross linked polystyrene resin (capacity 0.68 mmol/g) by solid-phase method, using Boc chemistry. The following side-chain protection groups were employed: benzyl (Ser, Thr), 2-chlorobenzoyloxycarbonyl (Lys), cyclohexyl (Asp), tosyl (Arg) and Np-benzoylohexymethyl (His) [22]. The peptides were cleaved from the resin by the HF procedure and lyophilized. The crude products were desalted on a Sephadex G-25 column (2.8 cm  $\times$  105 cm) in 30% CH<sub>3</sub>COOH and chromatographed twice by preparative RP-HPLC on a Vydac C-18 column (32 mm  $\times$  240 mm, 15–20 mm particle size). Separation was done with the use of a linear gradient of 0–15% acetonitrile in 0.1% trifluoroacetic acid. Relative molecular ions mass of the peptides were determined by fast atom bombardment mass spectrometry method (FAB

MS) on AMD-604 mass spectrometer. Amino-acid analysis was performed on a Beckman model 121M analyzer. The peptide was hydrolyzed with hydrochloric acid containing 1% of phenol at 110°C for 24 h.

**Table 1. Sequences of Tat model peptides**

HIV-1 Tat*	... <sup>39</sup> RKLGISYGRKKRRRQRRRAPQDSQTHQASLSKQPASQ <sup>76</sup> ...
Tat1 peptide	<sup>1</sup> GRKKRRRQRRRA <sup>11</sup>
Tat2 peptide	<sup>1</sup> GRKKRRRQRRRAPQDSQTHQASLSKQPA <sup>27</sup>

\*Data from [27].

### Isolation of RNA

Ribosomal 5S RNA was isolated from yellow lupin (*Lupinus luteus*) seeds by phenol extraction, purified on Sephadex G-75 and repurified on 15% polyacrylamide gel containing 7 M urea, 50 mM Tris/borate buffer, pH 8.3 and 1 mM EDTA (TBE), as described previously [23]. tRNA<sup>Phe</sup> extracted from yeast was additionally purified on 15% polyacrylamide gel containing 7 M urea in 50 mM TBE buffer. 5S rRNA and tRNA were 3' end labelled with [<sup>32</sup>P]pCp and RNA ligase [24]. <sup>32</sup>P-Labelled RNAs were purified by 10% polyacrylamide gel electrophoresis (PAGE) in the presence of 7 M urea, eluted from the gel and renatured [25].

### RNA binding reactions and electrophoretic mobility gel shift assays

The RNA-Tat peptide complex formation assay was performed at 22°C for 40 min in a buffer containing 50 mM Tris/HCl, pH 7.5, 70 mM NaCl, 1 mM EDTA, 0.1% Nonidet P-40, 1 nM 5S rRNA (or tRNA<sup>Phe</sup>), 2 mg crude tRNA, 3 nM Tat peptide in a total volume of 10 ml. Electrophoretic analysis of the complexes was carried out on 0.7% agarose gel in 50 mM Tris/borate/EDTA (TBE) buffer.

### RNase footprint assay

For a footprint reaction of the RNA-peptide complex the following amounts of RNases were used T1 ( $2 \times 10^{-4}$  U), V1 ( $6 \times 10^{-2}$  U), S1 (3 U) (Pharmacia). For localization of Tat1 binding site, 4 mg of cold 5S rRNA or tRNA

and 40000 c.p.m. of labelled 5S rRNA or tRNA<sup>Phe</sup>, respectively, were digested with 0.04 U of T1 RNase in a buffer containing: 20 mM CH<sub>3</sub>COONa, pH 4.5, 7 M urea, 1 mM EDTA and 0.05% xylene cyanol, and ana-

lyzed on 10% polyacrylamide gel in the presence of 7 M urea in 0.09 M TBE buffer.

### Circular dichroism measurements

CD spectra of Tat1 and Tat2 peptides as well as RNAs and their complexes were recorded at room temperature on a Jasco J-20 spectropolarimeter over an 190–310 nm interval using a 1.0 mm pathlength cell. RNA-Tat peptide 1:1 complexes were prepared in 100 mM Tris, pH 7.0, using concentrations of  $5.2 \times 10^{-6}$  M and  $3.3 \times 10^{-6}$  M for tRNA and 5S rRNA, respectively. Concentration of the Tat peptides used for CD spectra was  $2 \times 10^{-4}$  M. The molar ellipticity of Tat-RNA complexes and free peptides was calculated per nucleotide of RNA.

### Capillary electrophoresis analysis

Free zone capillary electrophoresis (FZ-CE) (Beckman P/ACE System 2100) was used for analysis (measured at 214 nm) of peptides purity and RNA-Tat complexes formation. The samples of peptides were separated in 0.1 M sodium phosphate buffer, pH 2.5, using an uncoated fused silica capillary (75 mm × 57 cm, 50 cm to the detector). Runs were done at 18 kV (140 μA) and 25°C. Samples were injected by pressure for 5 s. The tRNA<sup>Phe</sup>-Tat1 complex formation was analyzed in 25 mM sodium phosphate buffer, pH 7.0, at constant voltage of 30 kV (68 μA) and 25°C.

All solutions and samples were filtered through 0.22 μm membrane filters. Some properties of Tat fragments are summarized in Table 2.

**Table 2. Physicochemical properties of Tat peptides.**T<sub>m</sub>, migration time, R<sub>t</sub>, retention time.

Compound	FAB-MAS Relative molecular mass M+H <sup>+</sup>		FZ-CE T <sub>m</sub> [min]	RP-HPLC R <sub>t</sub> [min]	Amino-acid analysis
	calc.	found			
Tat1	C <sub>58</sub> H <sub>114</sub> N <sub>32</sub> O <sub>13</sub> +H <sup>+</sup>		8.54	8.8	Glu1.10(1), Gly1.05(1) Ala1.00(1), Lys1.78(2), Arg5.85(6)
	1468.8	1468.5			
Tat2	C <sub>129</sub> H <sub>227</sub> N <sub>55</sub> O <sub>39</sub> +H <sup>+</sup>		13.28	9.67	Glu5.21(5), Ser3.11(3), Lys3.00(3), Pro1.79(2), Arg5.92(6), Asp0.96(1), Gly0.89(1), Thr0.91(1), Ala3.00(3), Leu0.87(1), His0.92(1)
	3173.6	3173.8			


**RESULTS AND DISCUSSION**

Many recent studies have been focused on determination of the mechanism of protein-nucleic acids recognition. The NMR data of the HIV TAR-Tat complex indicated that an arginine residue recognizes an UCU bulge [17]. This mode of interactions is totally different from those observed in some DNA-protein complexes, where direct binding of guanine and arginine residues was observed [26]. Why the interactions within RNA-protein and DNA-protein complexes are different and why arginine residue does not interact with guanine-rich loop? To solve this question we turned to native RNA molecules

for which there is a lot of structural data including their three dimensional structure, i.e. yeast tRNA<sup>Phe</sup> and lupin 5S rRNA.

tRNA<sup>Phe</sup> from yeast was found to contain a (5′)-UGGG sequence in the dihydrouridine loop. The tetranucleotide string resembles very well a part of the TAR RNA hairpin tip (Fig. 1). Two guanosine residues (G18G19), are conserved in all cytoplasmic tRNAs.


To check the specificity of RNA binding with the arginine rich peptide, we prepared two synthetic peptides, Tat1 and Tat2, 11 and 27 amino acid long, respectively (Table 1). As one can see, the amino-acid sequence of these peptides corresponds exactly with the RNA binding motif of the HIV-1 Tat protein.



**Figure 1.** The secondary structure of: A, TAR RNA of HIV-1 [17], B, tRNA<sup>Phe</sup> of yeast [20], C, 5S rRNA of *Lupinus luteus* [23]. UGGG sequence is highlighted by black box.

The first step in our studies was to check whether, or not, these peptides form complexes with tRNA<sup>Phe</sup>. Results of the agarose gel shift assay show that both Tat1 and Tat2 bind to yeast tRNA<sup>Phe</sup> (Fig. 2). In the presence of a three fold excess of the peptides in the reaction mixture, 100% of RNA was bound. The protein-RNA complex formation was also confirmed by capillary electrophoresis (Fig. 3). The data presented in Fig. 3 for Tat 1 indicate that the peaks corresponding to the Tat1 peptide and tRNA<sup>Phe</sup> with migration time ( $M_t$ ) of 3.57 min and 17.10 min, respectively, disappear after complex formation and a new peak corresponding to the Tat 1-tRNA<sup>Phe</sup> complex is the only one to be seen.


In order to determine the peptide binding site on the RNA studied, a limited RNase hydrolysis with RNase T1 (G-specific), S1 (single-stranded RNA specific), V1 (double-stranded RNA specific) was carried out. For these studies we used the Tat1 peptide at two different concentrations (Figs. 4 and 5). The footprints of the RNA-Tat2 complex were so smeared as to make their interpretation practically impossible. Therefore, further footprint analysis in this paper concerns the Tat1-RNA complexes. T1 RNase hydrolysis of tRNA<sup>Phe</sup> gave two weak bands corresponding to cuts at nucleotides G18 and G19 (Fig. 4). These cuts are absent in the digest of the complex. At the same time, there are no



**Figure 2.** Electrophoretic agarose gel-shift analysis of binding of Tat peptides to [3'-<sup>32</sup>P]5S rRNA (lanes 1-3) and [3'-<sup>32</sup>P]tRNA<sup>Phe</sup> (lanes 4-6) on 0.7% agarose gel (lanes 2, 5 — Tat1, 3, 6 — Tat2).

Arrow points the RNA-Tat peptide complexes.

differences in the hydrolysis by RNase V1 of tRNA alone and in the complex. Interestingly, RNase S1, in contrast to T1 RNase, hydrolyses tRNA<sup>Phe</sup> complexed with Tat1 at each of the three nucleotides G17G18G19 (Fig. 6A) but not the free tRNA. Results of




**Figure 3.** Free zone capillary electrophoresis of Tat1, tRNA<sup>Phe</sup> and Tat1-tRNA<sup>Phe</sup> complex.

Analysis were carried out in uncoated fused-silica capillary (75  $\mu$ m  $\times$  57 cm, 50 cm to the detector), constant voltage of 30 kV (68  $\mu$ A), temp. 25°C.

hydrolysis of the complex with T1, S1 and V1 RNases clearly suggest that the Tat1 peptide interacts with dihydrouridine loop of tRNA<sup>Phe</sup> molecule.


From the crystallographic structure of yeast tRNA<sup>Phe</sup> it is known that G18 and G19 of the loop D and C56 and Y55 in ribothymidine loop (T) form hydrogen bonds, respectively [20]. In the case of a single-stranded RNA specific enzyme (RNase S1), lack of cleavages at G18 and G19 in the absence of Tat1 and strong hydrolysis in its presence suggest that binding of the peptide induces in tRNA<sup>Phe</sup> some conformational changes so that the sugar-phosphate backbone in the complex becomes accessible. The guanosine residues in dihydrouridine loop are protected against T1 RNase by the peptide.

Circular dichroism spectra of free tRNA<sup>Phe</sup> (Fig. 7A) show a Cotton effect at 266 nm which disappears after addition of the Tat1 and Tat2 peptides; this means that the conformation of tRNA<sup>Phe</sup> becomes changed upon peptide binding. These changes may be interpreted as a disruption of the tertiary interaction between the D and T loops induced by these peptides [28, 29]. From other studies it is known that if the A-form of RNA is perturbed, the intensity of 265 nm band decreases and that of the other band at 240 nm increases [29]. These observations led us to the conclusion that the tertiary hydrogen bonds of tRNA, G19-C56 and G18-Y55, are disrupted in the complex of tRNA<sup>Phe</sup> with Tat peptides. Therefore, we postulate that the Tat1 and Tat2 peptides should form hy-



**Figure 4.** Autoradiogram of 10% polyacrylamide gel containing 7 M urea showing the products of [3'-<sup>32</sup>P]tRNA<sup>Phe</sup>-Tat1 complex hydrolysis by RNase T1 (lanes 4-6), RNase VI (lanes 7-9) and RNase S1 (lanes 10-12).

Lanes: 1, control: 5S rRNA incubated in the reaction buffer (10 min/22°C); 2, ladder; 4, 7, 10, tRNA<sup>Phe</sup>; 5, 8, 11, tRNA<sup>Phe</sup> + 1.45 nM Tat1 peptide; 6, 9, 12, tRNA<sup>Phe</sup> + 3 nM Tat1 peptide. Differences in hydrolysis pattern are indicated by arrow. Numbers on the left side correspond to nucleotides. Lane 3, T1 nuclease digestions for guanosine ladder.




**Figure 5.** Autoradiogram of 10% polyacrylamide gel containing 7 M urea of the products of [ $3'$ - $^{32}$ P]5S rRNA-Tat1 hydrolysis by RNase T1 (lanes 4-6), RNase V1 (lanes 7-9) and RNase S1 (lanes 10-12).

Lanes: 1, control: 5S rRNA incubated in reaction buffer (10 min/22°C); 2, ladder; 3, T1 nuclease digestions for guanosine ladder; 4, 7, 10, 5S rRNA; 5, 8, 11, 5S rRNA + 1.45 nM Tat1-peptide; 6, 9, 12, 5S rRNA + 3 nM Tat1-peptide. Differences in hydrolysis pattern are indicated by arrow. Numbers on the left side correspond to nucleotides.

drogen bonds with tRNA<sup>Phe</sup> in a similar way as zinc fingers of Zif268 in complex with DNA, e.g. mostly guanine-arginine [26].


Similar results were obtained from experiments on the interaction of Tat-peptides with 5S rRNA from yellow lupin, bearing the same nucleotide sequence, UGGG, as tRNA<sup>Phe</sup>. The agarose gel shift assay indicated that 5S rRNA also forms complexes both with peptide Tat1 and Tat2 (Fig. 2). The patterns obtained on hydrolysis of 5S rRNA-Tat1 com-

plex with RNases T1 and V1 (Fig. 5) show that nucleotides G89, G56 and 93-95 were not affected at all. However, strong bands due to digestion of the backbone at G85-G87 nucleotides by RNase S1 were observed (Figs. 5, 6A). This suggests that the Tat1 peptide interacts with D-loop and a part of stem IV of 5S rRNA molecule. According to the three-dimensional model of plant 5S rRNA, the nucleotides G85-G87 of the loop D are involved in interaction with C34-C36



**Figure 6.** The secondary structure of tRNA<sup>Phe</sup> from yeast (A) and 5S rRNA from *L. luteus* (B) with marked results of footprint experiments on the complex with Tat1-peptide.

Open arrows indicate the nucleotides protected from hydrolysis by RNase T1 and black ones the sites of enhanced digestion by RNase S1. Nucleotides protected from RNase V1 are in box.



of the loop C [21, 30]. The results of hydrolysis of the 5S rRNA-Tat complex (summarized in Fig. 6B) with RNases are similar to those observed with the tRNA<sup>Phe</sup>-Tat1 complex. Inspection of CD spectra (Fig. 7B) shows that maximum of Cotton effect of 5S rRNA at 265 nm decreases in the presence of Tat peptides, as it has been observed in the case of tRNA<sup>Phe</sup> (Fig. 7A). Interestingly, CD spectra of both RNA's complexed with Tat2 suggest that this peptide has a smaller influence on the conformation of either of them than Tat1. This observation is similar to the data obtained on interaction of poly dGm<sup>5</sup>dC with two synthetic peptides containing the KPKKKKEK sequence, 22 and 8 amino acids long, derived from lupin glutamyl-tRNA synthetase [31]. It was observed by CD that the presence of the shorter peptide brought about changes in B-DNA conformation of poly dGm<sup>5</sup>dC while the longer peptide did not produce such changes. These data suggest that shorter peptide has a higher ability to induce conformational changes in RNA,

**Figure 7.** CD-spectra of complexes of Tat1 and Tat2 with tRNA<sup>Phe</sup> (A) and 5S rRNA (B).

Full line, tRNA<sup>Phe</sup>, 5S rRNA; dashed line, Tat2-tRNA<sup>Phe</sup>, Tat1-5S rRNA; short dashed line, Tat1-tRNA<sup>Phe</sup>, Tat2-5S rRNA.



which can be explained by its flexible conformation. Indeed, NMR analysis of 25 amino acids long Tat-peptide, comprising the TAR binding domain of HIV-1, showed that in aqueous solution at pH 7.0 it has  $\alpha$ -helical structure [19]. This suggest that Tat2, due to its more stable structure, could not recognize RNA as precisely as the shorter Tat1. This conclusion is also supported by NMR analysis of arginine amide-DNA-hairpin complex. The 24-mer DNA with sequence: GATCGA-AACGTAGCGCCTTCGATC undergoes an adaptive conformational transition after binding of amino acid and finally arginine is encapsulated by the loop folded down toward the stem [31].

In summary, the data on Tat-RNA complexes presented in this paper lead to conclusion that the interactions between nucleotides C34C35C36 of loop C and G85G86G87 of loop D in the model of plant 5S rRNA structure [30] are very similar to those observed between the loops D and T in tRNA<sup>Phe</sup> from yeast. Another conclusion is that the UGGG sequence in the loop of RNA is a target for binding of arginine rich proteins. Finally, we postulate that the binding site for Tat protein is the guanosine rich loop and not the UCU bulge.

## REFERENCES

1. Freemont, P.S., Lane, A.N. & Sanderson, M.R. (1991) Structural aspects of protein DNA recognition. *Biochem. J.* **278**, 1–23.
2. Otwinowski, Z., Schevitz, R.V., Zhang, R.G., Lawson, C.L., Joachimiak, A., Marmorstein, R.Q., Luisi, B.F. & Sigler, P.B. (1988) Crystal structure of trp repressor/operator complex at atomic resolution. *Nature (London)* **335**, 321–326.
3. Suzuki, M. & Yagi, N. (1994) DNA recognition code of transcription factor in the helix-turn-helix, probe helix, hormone receptor and zinc finger families. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 12357–12361.
4. Choo, Y. & Klug, A. (1994) Toward a code for the interactions of zinc fingers with DNA: Selection of randomized fingers displayed on phage. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 11163–11167.
5. Greisman, H.A. & Pabo, C.O. (1997) A general strategy for selecting high affinity zinc finger proteins for diverse DNA target sites. *Science* **275**, 657–661.
6. Mattaj, I.W. (1993) RNA-recognition: A family matter. *Cell* **73**, 837–840.
7. Rould, M.A., Perona, J.J., Soll, D. & Steitz, T.A. (1989) Structure of *E. coli* glutamyl-tRNA synthetase complexed with tRNA<sup>Gln</sup> and ATP at 2.8 Å resolution. *Science* **246**, 1136–1142.
8. Ruff, M., Krishnavarmy, S., Boeglin, M., Poterszman, A., Mitschler, A., Podjarny, A., Rees, B., Thierry, J.C. & Moras, D. (1991) Class II aminoacyl transfer RNA synthetases: crystal structure of yeast aspartyl-tRNA synthetase complexed with tRNA<sup>Asp</sup>. *Science* **252**, 1682–1689.
9. Biou, V., Yaremchuk, A., Tukalo, M. & Cusack, S. (1994) The 2.9 Å crystal structure of *T. thermophilus* seryl-tRNA synthetase complexed with tRNA<sup>Ser</sup>. *Science* **263**, 1404–1410.
10. Valegard, K., Murray, J.B., Stockley, P.G., Stonehouse, N.J. & Liljas, L. (1994) Crystal structure of an RNA bacteriophage coat protein-operator complex. *Nature (London)* **371**, 623–626.
11. Oubridge, C., Ito, N., Evans, P.R., Teo, C.H. & Nagai, K. (1994) Crystal structure at 1.92 Å resolution of the RNA-binding domain of the U1A splicesomal protein complexed with an RNA hairpin. *Nature (London)* **372**, 432–438.
12. Burd, C.G. & Dreyfuss, G. (1994) Conserved structures and diversity of function of RNA-binding proteins. *Science* **265**, 615–621.
13. Łaziński, D., Grządzińska, E. & Das, A. (1989) Sequence specific recognition of RNA hairpins by bacteriophage antiterminators requires a conserved arginine rich motif. *Cell* **59**, 207–218.
14. Chen, L. & Frankel, A.D. (1994) An RNA-binding peptide from bovine immunodeficiency virus Tat protein recognizes an un-

- usual RNA structure. *Biochemistry* **33**, 2708–2715.
15. Weeks, K.M., Ampe, Ch., Schultz, A.S., Steitz, T.A. & Crothers, D.M. (1990) Fragments of the HIV-1 Tat protein specifically bind TAR RNA. *Science* **249**, 1281–1285.
  16. Puglisi, D., Tan, R., Calnan, B.J., Frankel, A.D. & Williamson, J.R. (1992) Conformation of the TAR RNA arginine complex by NMR spectroscopy. *Science* **257**, 76–80.
  17. Puglisi, D., Chen, L., Frankel, A.D. & Williamson, J.R. (1993) Role of RNA structure in arginine recognition of TAR RNA. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 3680–3684.
  18. Aboul-ela, F., Karn, J. & Varani, G. (1995) The structure of the human immunodeficiency virus type-1 TAR RNA reveals principles of RNA recognition by Tat-protein. *J. Mol. Biol.* **253**, 313–332.
  19. Mujeeb, A., Parslow, T.G., Yuan, Y.-Ch. & James, T.L. (1996) Aqueous solution structure of a hybrid Lentiviral Tat peptide and a model of its interaction with HIV-1 TAR RNA. *J. Biomolec. Struct. Dynamics* **13**, 649–660.
  20. Kim, S.H., Suddath, F.L., Quingley, G.J., McPherson, A., Sussman, J.L., Wang, A.H., J., Seeman, N.C. & Rich, A. (1974) Three dimensional tertiary structure of yeast phenylalanine transfer RNA. *Science* **185**, 435–440.
  21. Joachimiak, A., Nalaskowska, M., Barciszewska, M., Mashkova, T.D. & Barciszewski, J. (1990) Higher plant 5S rRNAs share common secondary structure. A new three domains model. *Int. J. Biol. Macromol.* **12**, 321–327.
  22. Horiki, K., Igano, K. & Inoue, K. (1978) Amino acids and peptides. Part 6. Synthesis of the Marrifield resin esters of N-protected amino acid of hydrogen bonding. *Chem. Lett.* 165–168.
  23. Barciszewska, M., Mashkova, T.D., Zwierzynski, T., Kisselev, L.L. & Barciszewski, J. (1986) The primary structure of yellow lupin seeds 5S ribosomal RNA. *Bull. Acad. Polon. Sci.* **34**, 369–373.
  24. Barciszewska, M., Dirheimer, G. & Keith, G. (1983) The nucleotide sequence of methionine elongator tRNA from wheat germ. *Biochem. Biophys. Res. Commun.* **114**, 1161–1168.
  25. Pieler, T. & Erdmann, V.A. (1982) Three dimensional structural model of eubacterial 5S RNA that has functional implications. *Proc. Natl. Acad. Sci. U.S.A.* **79**, 4599–4603.
  26. Pavletich, N.P. & Pabo, C.O. (1991) Zinc finger — DNA recognition: Crystal structure of a Zif268-DNA complex at 2.1 Å. *Science* **252**, 809–817.
  27. Loret, E.P., Vives, E., Ho Shing, P., Rochat, H., Rietschoten, J.V. & Johnson, W.C., Jr. (1991) Activating region of HIV-1 Tat protein: Vacuum UV circular dichroism and energy minimization. *Biochemistry* **30**, 6013–6023.
  28. Tan, R. & Frankel, A.D. (1992) Circular dichroism studies suggest that TAT RNA changes conformation upon specific binding of arginine to guanidine. *Biochemistry* **31**, 10288–10294.
  29. Loret, E.P., Georgel, P., Johnson, W.C., Jr. & Shing Ho, P. (1992) Circular dichroism and molecular modeling yield a structure for the complex of human immunodeficiency virus type 1 trans-activation response RNA and the binding region of Tat, the trans-acting transcriptional activator. *Proc. Natl. Acad. Sci. U.S.A.* **89**, 9734–9738.
  30. Barciszewska, M.Z., Erdmann, V.A. & Barciszewski, J. (1995) Ribosomal 5S RNA: Tertiary structure and interactions with proteins. *Biol. Rev. (Cambridge)* **71**, 1–25.
  31. Lin, H.C. & Patel, D.J. (1996) Encapsulating an amino acid in a DNA fold. *Nature Struct. Biol.* **3**, 1046–1060.