

Erratum

ls:

Page 38, Materials and Methods, Patient Samples

40 AML patients and 40 healthy controls were recruited from Ningbo First Hospital. Among the AML patients, male: 26, female:14; age: 35.21±6.14; physical activity state score, 0-1: 27, 2-4: 13; Type, M1: 3; M2: 5; M3: 11; M4: 13; M5: 6; M6: 2. The peripheral blood was collected and stored in liquid nitrogen immediately until needed. The patients who received chemotherapy or radiotherapy were excluded from this study. This study was approved by the Research Ethics Committee of Ningbo First Hospital20190126, January 26th, 2019, and each patient has signed an in formed consent form.

Should be:

Page 38, Materials and Methods, Patient Samples

40 AML patients and 40 healthy controls were recruited from Shaoxing People's Hospital. Among the AML patients, male: 26, female:14; age: 35.21±6.14; physical activity state score, 0-1: 27, 2-4: 13; Type, M1: 3; M2: 5; M3: 11; M4: 13; M5: 6; M6: 2. The peripheral blood was collected and stored in liquid nitrogen immediately until needed. The patients who received chemotherapy or radiotherapy were excluded from this study. This study was approved by the Ethics Committee of Shaoxing People's Hospital and each patient has signed an informed consent form.